INDEX TO VOLUME 55

New names and the final members of new combinations are in **bold face** type.

Abronia glabrifolia 76
Acacia 491
Acanthambrosia 185, 187, 189, 191; Bryantii 184
Acer negundo 109; rubrum 150
Achillea Millefolium 205, 208, 209, 211
Aesculus Hippocastanum 150
Agaricus melleus 513
Ailanthus 375
Alchemilla 508
Ambrosia 187, 189, 190, 195; aptera 184; artemisiifolia 205, 208, 209, 211; bidentata 184; cumanensis 184; elatior 184; peruviana 184; psilostachya 184, 185, 197; tenuifolia 184; trifida 184
Ambrosiaceae, Pollen grains in the identification and classification of plants—I. The 181
American Botanical Literature, Index to 77, 133, 171, 214, 256, 319, 421
Anaphalis margaritacea 205, 208, 211
Anemone 447
Angiosperms, The phylogeny of the 479
Antennaria Brainerdii 101; neodioica 204, 208, 211; plantaginifolia 204, 208, 211
Anthemis Cotula 205, 208, 211; nobilis 193, 194, 197
Antholithes ranaliformis 442, 446
Arctium minus 205, 208, 211
Artemisia 183, 190, 191, 197
ASHE, W. W., Notes on southeastern woody plants 463
Aspergillus niger 281
Asplenium calopteris 444; nigrum 307
Aster 210, 435–437, 439; cordifolius 204, 208, 211; divaricatus 204, 207, 208, 211; ericoides 204, 208, 211; laevis 204, 208, 211; lateriflorus 204, 208, 211; Lowrieanus 204, 208, 211; macrophyllus 204, 208, 211; novae-angliae 204, 208, 211; paniculatus 204, 208, 211; prenanthoides 204, 208, 211; puniceus 204, 208, 211; sagittifolius 204, 208, 211; umbellatus 204, 208, 211; undulatus 204, 208, 211
Astragalus aboriginum 120, aboriginum fastigiosum 122, 123; alpinus 119, 130–132; alpinus Brunetianus 125, 131; alpinus giganteus 131; altus 161; andinus 130; Blakei 125, 128, 131; cobrensis 162; Cottoni 121; elegans 128, 130; eucosmus 126, 129; flexuosus 130; Forwoodii 122; frigidus 131; Funstoni 127; giganteus 131; glabriusculus 120; glabriusculus major 122; Gormani 132; guatemalensis Lozani 161; guatemalensis oaxacanus 160; Hallii 130; Harringtonii 127, 128; Hartwegii 163, 164; Jesupi 125, 128, 131; labradoricus 124, 125, 126, 128, 131; leptocarpus 157, 158; linifolius 75; Macounii 125, 128, 131; occidentalis 125, 126, 131; olympicus 122; oroboides 128, 132; Painteri 156; platytropus 130; Pueblae 164; Purpusi 164; rafaelensis 75; recurvus 162; reptans 159, 160; Robbinsii 125, 129, 131; Robbinsii borealis 125; Robbinsii Jesupi 125; Robbinsii occidentalis 126, 128; Rosei 163; Rusbyi 157, 162, 163; Rusbyi longissimus 162; Seatoni 164; secundus 125, 131; spatiosus 122; stenophyllus 122; straturensis 162; strigulosus 130, 156–160, 163; strigulosus brevidentatus 156, 162; strigulosus gracilis 159, 160, 161; tolucanus 156; vaginatus 121; Williamsii 132
Atelophragma aborignum 119–122; **alpiniforme** 128, 129; **alpinum** 129–131; **altum** 156, 161; **Blakei** 124, 125; **bracteatum** 126, 161; Brandegei 128, 129; **cobrense** 156, 162; **Collieri** 124, 127; Cottoni 120, 121; elegans 126, 128, 129; **Fernaldi** 124, 126; Forwoodii 120, 122, 123; glabriusculum 120, 122, 123; **guatemalense** 155, 159, 161; **Harringtonii** 124, 126; **Harshbergeri Herriotii** 120, 123; **hidalgense** 155, 159; **jaliscense** 155, 157; **Jesupi** 124, **labradoricum** 130, 131; lineare 120, 123; **longissimum** 156, 162; **Lozani** 156, 158, 161; Macounii 124, 126, 128; **oaxacanum** 155, 160; occidentale 128, 130; oroboides 119, 132; **Painerti** 155, 156; **potosinum** 156,

160, 161; **Pueblae** 163; 164, **Purpusi** 164; **reptans** 155, 159; Richardsonii 121; **Robbinsii** 124, 128; **Rosei** 156, 163; **Rusbyi** 156, 162; **Seatoni** 164; Shearii 128, 130; **straturense** 156, 162; **strigulosum** 155, 157–161, 163; **tioides toluccanum** 155, 156; **Townsendii** 156, 163; wallowense 120, 122; **Williamsii** 132; **zacatecanum** 155, 157
Atelophragma, Notes on Fabaceae—X, 119; –XI, 155
Atlantic coastal plain-XV, Contributions to the Mesozoic flora of the 441
Aucuba 308, 309
BALDWIN, HENRY I. A comparison of the available moisture in sod and open soil by the soil-point method 251
Bamboo, A study of the grand period of growth in 327
Bambusa 334; arundinacea 331; baccifera 331; Balcoa 331; gigantea 331, 332; macroculmis 332, 366; spinosa 334, 335; Tulda 332; verticillata 332; vulgaris 332, 333, 335, 362, 366, 371, 374; vulgaris vittata 332, 366
Barnadesia 449, 454–456, 458–462; berberoides 451, 462; divaricata 452, 461; glomerata 451; hirsuta 451; inermis 451–453, 461, 462; macrocephala 451; odorata 451; polyacantha 451; rosea 451; spinosa 451, 455; Trianae 450, 451, 457, 462; venosa 450, 451, 462
Barnadesia, Pollen grains in the identification and classification of plants—II. 449
Begonia semperflorens 311
Berlandiera lyrata 192
BERRY, EDWARD W. Contributions to the Mesozoic flora of the Atlantic coastal plain—XV 441
Bidens 210; cernua 205, 208, 211; comosa 205, 208, 211; connata 192; frondosa 205, 208, 211; laevis 192; vulgata 192, 205, 208, 211
BOLENBAUGH, ALTA. Microsporogenesis in Tropaeolum majus with special reference to the cleavage process in tetrad formation 105
Bolivia, New species of Peperomia from 169
Botanical Literature, Index to American 77, 133, 171, 214, 256, 319, 421, 467, 554
Botrytis cinerea 329
Brainerd, Ezra (1844–1924) (*portrait*) 91
Brassica oleracea 223, 545
Bromus inermis 380
Brousonettia 375
Bryophyllum 328; calycinum 311
Burgess (1855–1928) (*portrait*), Edward Sandford 433
BUSH, BENJAMIN FRANKLIN. Quercus neo-Ashei 247
Callistemon 491
Callisteris violacea Greene 69
CAMPBELL, DOUGLAS HOUGHTON. The phylogeny of the Angiosperms 479
Canada, A revision of the Plantago patagonica group of the United States and 406
Canna 491
Cannabis 376
Capparis proeocenica 442
Carduus nutans 205, 208, 211
Carex 93, 95; Brainerdii 96
Carya alba 250; arkansana 248; Buckleyi 248; leiodermis 465; leiodermis callicoma 465
Casuarina 487
Caltha 447; palustris 12
Celastrophyllum Ripleyanum 442
Celtis Lindheimeri 248; reticulata 248
Centurea Jacea 205, 208, 210, 211; maculosa 205, 208, 211
Cercis Texana 248
Cestrum nocturnum 366
Chaetomium 516, 519, 523
Chamissomenia 462
Chara 14; coronata 17, 39; delicatula 20, 39
Characeae, Nuclear and cell division in the antheridial filaments of the 11
Chorisiva 189, 190, 193, 194, 198; nevadensis 184, 185, 196, 197
Chrysanthemum Leucanthemum 205, 208, 209, 211
Chuquiraga 461, 462; ferox 452, 453, 460; floribunda 452, 453, 459, 462; glabra 453; leptacantha 452, 453; vagans 453; varians 453
Cichorium Intybus 205, 208, 211
Cirsium arvense 205, 208, 211; lanceolatum 205, 208, 210, 211; muticum 205, 208, 211
Cobaea scandens 106
Coleus Blumei 311
Colorado, New plants from 75
Comparison of the available moisture in sod and open soil by the soil-point method, A 251

Conocephalum 9; conicum 7
Contributions to the Mesozoic flora of the Atlantic coastal plain—XV 441
Convolvulus lupuliformis 499; sepium 499, 500, 508
Coprinus stercorarius 513
Coreopsis major 192; tinctoria 192
Corsinia 3
Corydalis cava 12; pumila 12
Cosmos 374; bipinnatus 192
Costus Mooreana 165
Crataegus aestivalis **Dormonae** 464; Crus-galli 98; Pringlei 98
Cremanium rubens 118; trinitatis 118
Crepis 105, 110
Crocus 508
Crossostephium artemisioides 191
Ctenophyllum linifolium 75
Cucurbita 107; maxima 105; Pepo 371
Cuscuta Epithymum 499, 500, 508; Gronovii 499, 500, 508; lupuliformis 499
Cyclachaena 189, 190, 197; ambrosiaefolia 184, 185, 186, 197; lobata 184; pedicellata 184; xanthifolia 184, 186
Cystium 130
Cytological study of Ipomoea trifida, A morphological and 499
Dahlia pinnata 192
DAYTON, WILLIAM A. A new Gilia from the Montezuma National Forest 73; Callisteris violacea Greene 69
Dendrocalamus 333, 334, 361, 366, 367, 369, 371; giganteus 333–335, 359, 360, 367
Diapensia 94
Dichogamy in flowering plants 141
Dicorea 186, 189, 190, 197; Brandegei 184; canescens 184
Dictamnus albus 21
Didymosorus comptoniifolius 444
Diodia 508
Dioon 12
Dryopteris 312; normalis 312, 314
Edward Sandford Burgess (1885–1928) (*portrait*) 483
EGGLESTON, WILLARD W. Ezra Brainerd (1844–1924) (*portrait*) 91
EISENMENGER, WALTER S. Toxicity, additive effects, and antagonism of salt solutions as indicated by growth of wheat roots 261
Ephedra 488
Epilobium hirsutum 306
Equisetum 265
Erechtites 210; hieracifolia 205, 208, 211
Erigeron annuus 204, 208, 211; canadensis 204, 208, 211; philadelphicus 204, 208, 211; ramosus 204, 208, 211
Erysiphe 514; graminis 41, 44, 47, 65
Erythrinolepis 448
Eucalyptus 491
Euonymus 308
Eupatorium maculatum 204, 208, 211; urticaefolium 204, 208, 211
Euphorbia Lathyris 12
Euphrosyne 186, 189, 190, 197; parthenifolia 184
Ezra Brainerd (1844–1924) (*portrait*) 91
Fabaceae—X, Atelophragma, Notes on 119; —XI. 155
FARR, CLIFFORD H. Studies on the growth of root hairs in solutions—VII. Further investigations on collards in calcium hydroxide 223; —VIII. Structural and intracellular features of collards in calcium nitrate 529
Ficus 483
Flotovia 453
Franseria 185, 189, 195; acanthicarpa 184; albicaulis 184; bipinnatifida 184; deltoidea 184, 187; dumosa 184; ilicifolia 184, 187; tenuifolia 184, 187
Fritillaria 14; imperialis 11, 21
Fucus 13
Fuligo 13
Fumago 513
Further investigations on collards in calcium hydroxide, Studies on the growth of root hairs in solutions—VII. 223
Galinsoga parviflora 192
Gigantochloa 334; aspera 334; levis 335
Gilia aggregata 70, 71; pulchella 71; violacea 71
Gilia from the Montezuma National Forest, A new 73
Gladiolus 508
GLEASON, H. A. Miconia minutiflora and allied species 117
Gleichenia 444, 445; delicatula 442, 443
Glomerella cingulata 268
Glossocentrum collinum 117
Gnaphalium decurrens 205, 208, 211; uliginosum 205, 207, 208, 211
Gnetum 492

Griffithsia 13
Grindelia squarrosa 204, 208, 211
Guardiola angustifolia 192; platyphylla 192
Gunnera 491
Hamamelis virginiana 105
Hamosa 158
Haplostephium Passerina 458
HEIN, ILLO. Studies on morphogenesis in fungous mycelia 513
Helianthus 376, 380; annuus 192, 194, 195, 197; divaricatus 205, 208, 211; strumosus 192, 205, 208, 211
Hemerocallis 508
Hemizonia citroidora 192; scabrella 192
Hicoria leiodermis callicoma 465; leiodermis **mollisima** 465; mollissima 465
Hieracium aurantiacum 205, 208, 211
Hordeum vulgare 371
HOWE, MARSHALL AVERY. Edward Sandford Burgess (1855–1928) (*portrait*) 433
Hydrocharis 546
Hydrodictyon 13
Hymenoclea 187, 189, 191; fasciculata 184; monogyra 184; Salsola 184
Hypericum calycinum 501
Impatiens sultani 311
Index to American botanical literature 77, 133, 171, 214, 256, 319, 421, 467, 554
Inula Helenium 205, 208, 211
Ipomoea purpurea 499; trifida 499–510; trifida Torreyana 500
Ipomoea trifida, A morphological and cytological study of 499
Isoetes 494
Iva 183, 185, 186, 189–191, 197; angustifolia 184; axillaris 184, 197; cheiranthifolia 184; ciliata 184; frutescens 184; Hayesiana 184; oraria 184
Juglans cinera 145; regia 145
Juniperus monosperma 248
KARLING, JOHN S. Nuclear and cell division in the antheridial filaments of the Characeae 11
KENYAN, FRANCES MARION GREENE. A morphological and cytological study of Ipomea trifida 499
Lactuca 106; canadensis 205, 208, 211; scariola integrata 205, 208, 211; spicata 205, 208, 211
Leuciva 189, 190, 197; dealbata 184
Lilium 142, 491; giganteum 494
Limnobium 546
Long time experiments with plants in closed containers 305
Lupinus albus 265
Lychnophora rosmarinifolia 458; uniflora 458
Lythrum Salicaria 306, 307
Madia sativa 192
Magnolia 105, 483, 495; acuminata ludoviciana 464; acuminata ozarkensis 464; virginiana **parva** 464
Marsilia 12
MCALLISTER, FREDERICK. Sex ratio and chromosomes in Riccia Curtisii 1
Medicago 508
Megalospora affinis 513
Melampodium aureum 192; camphoratum 192
Melastoma minutiflora 117
Melilotus alba 107, 111
Mentha 244
Mesembryanthemum 491
Mesozoic flora of the Atlantic coastal plain—XV, Contributions to the 421
Metrosideros 491
Miconia **borealis** 117, 118; glossocentra 117; micrantha 117; myriantha 117, 118; trinitatis 118
Miconia minutiflora and allied species 117
Microsporogenesis in Tropaeolum majus with special reference to the cleavage process in tetrad formation 105
Mimosa 311, 313; pudica 311, 312, 314
Mimusops balata Schomburghii 447; **Collinsi** 442, 447; duplicata 447; praenuntia 447
Monilia fimicola 519
Montezuma National Forest, A new Gilia from the 73
Moriconia americana 445; cyclotoxon 442, 445
Morphogenesis in fungous mycelia, Studies on 513
Morphological and cytological study of Ipomoea trifida, A 499
Mulfordia 165; **Boliviana** 166, 167
Mulfordia, A new genus of the Zingiberaceae 165
Muscites 443; **Lesquereuxi** 442
Mutisia campanulata 455; viciaefolia 454, 462
Mycelia, Studies on morphogenesis in fungous mycelia 513
Mycogone 519

Myrica ornata 442; Torreya 442; Wadii minor 442
National Forest, A new Gilia from the Montezuma 73
Nectandra prolifica 442
New genus of the Zingiberazceae, Mulfordia, A 165
New Gilia from the Montezuma National Forest, A 73
New plants from Colorado 75
New species of Peperomia from Bolivia 169
Nicotiana 106, 313; Tabacum 311–314
Nitella 14; flexilis 19, 38, 39
Notes on Fabaceae—X. Atelophragma 119;—XI. 155
Notes on Scrophulariaceae of the northwestern United States 315
Notes on southeastern woody plants 463
Nuclear and cell division in the antheridial filaments of the Characeae 11
Oenothera 108
Opuntia 306
OSTERHOUT, GEORGE E. New plants from Colarodo 75
Oxalis stricta 311
Oxymitra 3
Oxytenia 186, 187, 190, 193, 194, 197, 198, acerosa 184, 185
Pacourina edulis 457
Pandanus 491, 492, 495
Parnassia 105
Parthenice 183, 193, 195, 196; mollis 192, 194, 198
Parthenium 183, 193–196, 198; hispidum 192; Hysterophorus 192; incanum 192; integrifolium 192; tomentosum 192
Pedicularis 317; ornithorhynchos 317; rainierensis 317
Penicillium 518
PENNELL, FRANCIS W. Notes on Scrophulariaceae of the northwestern United States 315
Peperomia 491; **aceramarcana** 169; **nequejahuirana** 169; **okarana** 170; **pitiguayana** 169; **ticunhuayana** 170
Peperomia from Bolivia, New species of 169
Phaca alpina 130, 131; elegans 128; frigida 132; glabriusculum 123; quinqueflora 164; Robbinsii 124
Pharbitis purpurea 500
Phaseolus multiflourus 327, 371; vulgaris 106, 508
Phyllites hydrocharitoides 446
Phyllostachys aurea 332, 366, 367; bambusoides 335, 357, 360, 361; mitis 332, 333, 357, 360, 364, 366, 367, 373; nigra 327–401; quadrangularis 327–405; violascens 357; viridi-glaucescens 332, 366, 367
Phylogeny of the Angiosperms, The 479
Picea canadensis 329; glauca 252
Plantago 417, 419; argyrea 414; aristata 408-411, 413; aristata Nuttallii 409, 410; brunnea 418, 419; Californica 417, 418; curta 412; dura 417, 418; erecta 417, 418; fastigiata 419; filiformis 409, 410; gnaphalioides 412; gnaphalioides aristata 409; Gooddingii 406, 419; Helleri 408, 414-416; Hookeriana 415, 416; Hookeriana **californica** 408, 417; Hookeriana **nuda** 408, 416; Hookeriana **typica** 408, 415; ignota 411, 412; inflexa 415, 416; insularis 418; insularis fastigiata 409, 419, 420; insularis scariosa 419; insularis **typica** 409, 418; Lagopus 412; lanatifolia 415; minima 406, 419, 420; Nuttallii 409; oblonga 411, 412; obversa 417; patagonica 406, 407; patagonica aristata 406, 409; patagonica californica 417; patagonica gnaphalioides 406, 412; patagonica lanatifolia 406, 415; patagonica nuda 406, 416, 417; patagonica spinulosa 406, 411; picta 412, 414; Purshii 407, 412, 414, 416; Purshii **argyrea** 408, 414; Purshii **typica** 408, 411–415; scariosa 419, 420; speciosa 417, 418; spinulosa 410–412, 414; spinulosa oblonga 408, 411, 413; spinulosa **typica** 408, 410, 411; squarrosa 409; tetrantha 417, 418; verticillata 411; Wrightiana 416; xerodea 412, 414
Plantago patagonica group of the United States and Canada, A revision of the 406
Plants, Dichogamy in flowering 141
Plants from Colorado, New 75
Plants in closed containers, Long time experiments with 305
Plants, Notes on southeastern woody 463
Plants—I. The Ambrosiaceae, Pollen grains in the identification and classification of 181
Platycodon grandiflorum 142, 144
Podophyllum 108

POE, IONE. A revision of the Plantago patagonica group of the United States and Canada 406
Polemonium caeruleum 95
Pollen grains in the identification and classification of plants—I. The Ambrosiaceae 181; —II Barnadesia 449
Polygonum Bistorta 12; Persicaria 306, 307
Polymnia canadensis 192; Uvedalia 192
Polystichum Braunii 94
Polytrichum commune 252
Populus 483
PORTERFIELD, WILLARD M., JR. A study of the grand period of growth in bamboo 327
Potamogeton 493; filiformis 446; foliosus 446; **hydrocharitoides** 442, 446; pectinatus 446; trichoides 446
Potentilla 501
Preissia commutata 433, 439; quadrata 434
Prenanthes 210; alba 205, 208, 211; altissima 205, 208, 211; trifoliolata 205, 208, 211
Puccinia glumarum 41
Pyronema 514
Quamoclidion cordifolium 75
Quercus annulata 248; Ashei 247; imbricaria 250; Laceyi 248; marilandica 247–250; marilandica Ashei 247, 248, 250; minima 465; Mohriana 248; **neo-Ashei** 247, 248, 250; **pygmea** 465; stellata 250; velutina 250; virginiana 248; virginiana dentata 465; virginiana pygmea 465
Ranunculus 447, 495
Raphaelia minuta 442, 443; neuropteroides 442, 443
Relation to the distribution of certain Compositae to the hydrogen-ion concentration of the soil 199
Revision of the Plantago patagonica group of the United States and Canada, A 406
Rhus aromatica 466; illinoiensis 465, 466; illinoiensis **formosa** 466; trilobata 466
Riccardia 9; pinguis 7
Riccia crystallina 9; fluitans 9; Frostii 9; glauca 9; lutescens 9; synspora 1
Riccia Curtisii, Sex ratio and chromosomes in 1
Romulea 508
Rosa 501
Rubus 102, 501
Rudbeckia hirta 192, 205, 208, 211; laciniata 192, 205, 208, 211
Ruppia 446
RUSBY, H. H. Mulfordia, a new genus of the Zingiberaceae 165
RYDBERG, PER AXEL. Notes on Fabaceae—X. Atelophragma 119; —XI. 155
Salix, 491; purpurea 148; viminalis 148
Salt solutions as indicated by growth of wheat roots, Toxicity, additive effects, and antagonism of 261
Salvia 374; sylvestris 76
Sarracenia 106
Scenedesmus quadricauda 308
Schlectendalia 453, 459–461; luzulaefolia 462
Schmaltzia formosa 465; illinoensis 465; serotina 465
Sclerotinia Libertiana 518
Scolymus hispanicus 456, 462
Scrophularia 315; californica 315, 316; lanceolata 316, 317; leporella 316; occidentalis 317; **oregana** 316
Scrophulariaceae of the northwestern United States, Notes on 315
Selaginella 12, 313, 443; Emiliana 312; laciniata 442
Seneceo aureus 205, 208, 211; **taraxacoides nudatus** 76; vulgaris 205, 208, 211
Sex ratio and chromosomes in Riccia Curtisisi 1
Silphium albiflorum 192; ovatifolium 192; perfoliatum 192
Soaresia velutina 458
Soil by the soil-point method, A Comparison of the available moisture in sod and open 251
Soil, Relation of the distribution of certain Compositae to the hydrogen-ion concentration of the 199
Solidago altissima 204, 208, 211; arguta 204, 208, 211; bicolor 204, 208, 211; caesia 204, 208, 211; canadensis Hargeri 204, 208, 211; graminifolia 204, 208, 211; juncea 204, 208, 211; latifolia 204, 208, 211; nemoralis 204, 208, 211; patula 204, 208, 211; rugosa 204, 208, 211; serotina 204, 208, 211; uniligulata 204, 208, 211
Sonchus asper 205, 208, 211
Sordaria 517, 519–521; fimiseda 516, 518
Sparganium 492, 493
Spathyema foetida 144

Sphaerocarpos californicus 1, 2; cristatus 1; Donnellii 1, 2, 6; terrestris 1, 2; texanus 2, 7
Sphaerotheca 514
Spilanthes americana 192; Beccabunga 192
Spiracantha cornuifolia 459
Spirogyra 264, 268, 281
Stilpnopappus ferrugineus 458
Stokesia 453, 457; laevis 462
STOUT, A. B. Dichogamy in flowering plants 141
Structural and intracellular features of collards in calcium nitrate, Studies on the growth of root hairs in solutions—VIII.
Studies on morphogenesis in fungous mycelia 513
Studies on the growth of root hairs in solutions—VII. Further investigations on collards in calcium hydroxide 223;—VIII. Structural and intracellular features of collards in calcium nitrate 529
Study of Ipomoea trifida, A morphological and cytological 499
Study of the grand period of growth in bamboo, A 327
Susceptbility of wheat to mildew as influenced by salt nutrition 41
Tanacetum vulgare 205, 208, 210, 211
Taraxacum 210; officinale 205, 208, 211
Thalictrum 491
Thallocarpus Curtisii 1
Thuja occidentalis 312
Tilia creno-serrata **acuminata** 465
Tium 119, 131, 158
Torenia asiatica 508, 509
Townsendia diversa 75; strigosa 76
Toxicity, additive effects, and antagonism of salt solutions as indicated by growth of wheat roots 261
Tradescantia 244
Tragopogon 110; pratensis 456
TRELEASE, HELEN M. See TRELEASE, SAM F.
TRELEASE, SAM F., and HELEN M. TRELEASE. Susceptibility of wheat mildew as influenced by salt nutrition 41
TRELEASE, WILLIAM. New species of Peperomia from Bolivia 169
Trianea 546
Trichia 13
Trillium 494
Triticum vulgare 371
Trollius 447
Tropaeolum 244
Tropaeolum majus with special reference to the cleavage process in tetrad formation, Microsporogenesis in 105
Tulipastrum acuminatum **ludovicianum** 464; acuminatum **ozarkense** 464
TURNER, J. ARTHUR. Relation of the distribution of certain Compositae to the hydrogen-ion concentration of the soil 199
Tussilago Farfara 205, 208, 210, 211
Typha 165
United States and Canada, A revision of the Plantago patagonica group of the 406
United States, Notes on Scrophulariaceae of the northwestern 315
Urtica pilulifera 12
Vaillantia 508
Vaucheria 13, 21, 26, 32; repens 22; terrestris 22; uncinata 22, 24; jucunda 459; stellaris 457; Wrightii 457
Viburnum bracteatum 464; molle 463, 464; **ozarkense** 463
Vicia Faba 105
Vinca 311; rosea alba 311
Viola 99, 102
WALLACE, RAYMOND H. Long time experiments with plants in closed containers 305
Wheat roots, Toxicity, additive effects, and antagonism of salt solutions as indicated by growth of 261
Wheat to mildew as influenced by salt Nutrition, Susceptibility of 41
Wielandiella 481, 490
Williansoniella 481
WODEHOUSE, R. P. Pollen grains in the identification and classification of plants.—I. The Ambrosiaceae 181;—II. Barnadesia 449
Woodwardia orientalis 312
Xanthium 183, 185, 187, 189, 195, 197; ambrosiaefolium 197; brevirostre 184; canadense 184; catharticum 197; chinense 184; globosum 184; pennsylvanicum 184, 197; speciosum 184; spinosum 184, 187, 197
Zannichellia 446
Zea Mays 371
Zingiberaceae, Mulfordia, A new genus of the 165
Zizyphus Ripleyensis 442